

Apparent violation of thermodynamics second law under Van der Waals gas expansion in vacuum

Yu.A.Koksharov

General Physics Department, Faculty of Physics, M.V.Lomonosov Moscow State University, Moscow, Russia, 119992

E-mail: koksharov@nm.ru

PACS numbers: 01.40.Fk;01.55.+b;05.70.-a

Submitted to: *EJP*

Abstract. Examining textbooks of thermodynamics we have not able to find a proof of increasing of entropy under free adiabatic expansion in vacuum based on the state equation of the Van der Waals gas. This communication remedies the situation. During proving we faced with an amusing example of apparent violation of thermodynamics second law.

The paradoxes and mistakes are not rare in textbooks of thermodynamics [1,2]. Many of them are concerned with entropy and the second law of thermodynamics [1-3]. An analysis of thermodynamics paradoxes is instructive for students and can be useful supplement to ordinary problems.

The second law is often deduced in classical thermodynamics as a principle of increasing of entropy in irreversible adiabatic processes [4]. For example, when a gas makes a free adiabatic expansion in vacuum its entropy S should increase. This is true for any gas regardless of its state equation due to the following general formula [5]:

$$\left. \frac{\partial S}{\partial V} \right|_U = \frac{p}{T} > 0,$$

where V is the system volume, U is the internal energy, T is the absolute temperature, p is the pressure.

In case of an ideal gas the increase of entropy in the process of free adiabatic expansion in vacuum can be shown plainly using the first law of thermodynamics

$$\delta Q = dU + \delta A, \quad (1)$$

where δQ is the heat received by the gas from its surroundings, δA is the work of external forces, which surroundings exert on the gas; the entropy definition

$$dS = \frac{\delta Q}{T}; \quad (2)$$

and the equation of state of an ideal gas

$$pV = \nu RT,$$

where R is the gas constant, ν is the mole number.

Indeed, in that case any infinitesimal change of S is equal to

$$dS = \frac{1}{T}(dU + \delta A) = \frac{1}{T}(C_V dT + p dV) = C_V \frac{dT}{T} + \nu R \frac{dV}{V}, \quad (3)$$

where C_V is the isovolumic specific heat.

The integration of equation (3) yields a change of the entropy when the ideal gas goes from an initial state 1 to a final state 2:

$$\Delta S = C_V \ln\left(\frac{T_2}{T_1}\right) + \nu R \ln\left(\frac{V_2}{V_1}\right) \quad (4)$$

Note that though equations (2,3) are valid only for reversible processes, equation (4) is applicable also to non-reversible processes, if initial and final states are thermal equilibrium.

We consider the adiabatic process, and, therefore, $Q = 0$. Besides, the total system volume $V_2 = \text{const}$, and we get $A = 0$. Hence, $\Delta U = Q - A = 0$ and for the ideal gas

$$\Delta T = \frac{\Delta U}{C_V} = 0 \rightarrow T_2 = T_1.$$

Using equation (2) and taking into account $V_2 > V_1$, we get

$$\Delta S = \nu R \ln\left(\frac{V_2}{V_1}\right) > 0.$$

A more complicate task is to develop a simple proof of entropy increasing under adiabatic expansion in vacuum in case of the Van der Waals gas. The state equation of the Van der Waals gas is written usually as [6]

$$\left(p + \frac{a}{V_\mu^2}\right)(V_\mu - b) = RT, \quad (5)$$

where $V_\mu = \frac{V}{\nu}$ is the molar volume, a, b are Van der Waals constants.

Using equations (1,5) we get formulas for changes of the entropy and the internal energy of the Van der Waals gas [7]:

$$\Delta S = C_{\mu,V} \ln\left(\frac{T_2}{T_1}\right) + R \ln\left(\frac{V_{\mu,2} - b}{V_{\mu,1} - b}\right) \quad (6)$$

$$\Delta U = C_{\mu,V}(T_2 - T_1) + a\left(\frac{1}{V_{\mu,1}} - \frac{1}{V_{\mu,2}}\right), \quad (7)$$

where $C_{\mu,V}$ is the isovolumic molar specific heat.

Since $\Delta U = 0$, it follows from equation (5):

$$T_2 = T_1 - \frac{a(V_{\mu,2} - V_{\mu,1})}{C_{\mu,V}V_{\mu,1}V_{\mu,2}} \quad (8)$$

Using $V_{\mu,2} > V_{\mu,1}$, from equation (8) we get $T_2 < T_1$, i.e., the Van der Waals gas always refrigerates under free adiabatic expansion in vacuum.

Substituting equation (8) for T_2 in equation (6), we obtain:

$$\Delta S = C_{\mu,V} \ln\left(1 - \frac{a(V_{\mu,2} - V_{\mu,1})}{C_{\mu,V}V_{\mu,1}V_{\mu,2}T_1}\right) + R \ln\left(\frac{V_{\mu,2} - b}{V_{\mu,1} - b}\right) \quad (9)$$

Taking into account $V_{\mu,2} > V_{\mu,1} > b$, we conclude that in equation (9) the second item is always positive, while the first item is always negative. One would think, that assuming the values of $a, b, V_{\mu,1}, V_{\mu,2}$ in equation (9) are fixed, we could choose T_1 sufficiently small, so that the sum (9) becomes negative (or equal to zero). However, **this conclusion conflicts with the second thermodynamics law**. Probably, our assumption is not correct. Let us prove that.

We begin with an inequality (see equation (6)):

$$\Delta S = C_V \ln\left(\frac{T_2}{T_1}\right) + R \ln\left(\frac{V_{\mu,2} - b}{V_{\mu,1} - b}\right) > 0 \quad (10)$$

From equation (10) it follows that:

$$\left(\frac{T_2}{T_1}\right)\left(\frac{V_{\mu,2} - b}{V_{\mu,1} - b}\right)^{\frac{R}{C_V}} > 1 \rightarrow$$

$$T_2(V_{\mu,2} - b)^{\frac{R}{C_V}} > T_1(V_{\mu,1} - b)^{\frac{R}{C_V}}. \quad (11)$$

Substituting T_2 from equation (8) in equation (11), we get:

$$T_1((V_{\mu,2} - b)^{\frac{R}{C_V}} - (V_{\mu,1} - b)^{\frac{R}{C_V}}) > (V_{\mu,2} - b)^{\frac{R}{C_V}} \frac{a}{C_V} \frac{V_{\mu,2} - V_{\mu,1}}{V_{\mu,1}V_{\mu,2}} \quad (12)$$

Assuming $V_{\mu,2} = V_{\mu,1} + dV$, where $dV \ll V_{\mu,1}$, from equation (12) we have:

$$\frac{RT}{C_V}(V_{\mu} - b)^{\frac{R}{C_V}-1}dV > (V_{\mu} - b)^{\frac{R}{C_V}} \frac{a}{C_V} \frac{dV}{V_{\mu}^2} \rightarrow$$

$$RT > a \frac{V_{\mu} - b}{V_{\mu}^2}, \quad (13)$$

where $V_{\mu} = V_{\mu,1}$, $T = T_1$.

However, from equation (5) it follows that

$$a \frac{V_{\mu} - b}{V_{\mu}^2} = RT - p(V_{\mu} - b) < RT, \quad (14)$$

when $p > 0$ and $V_{\mu} - b > 0$.

It remains to note that equations (13) and (14) are equivalent. Because all transformations from equation (10) to equation (13) are identical, we conclude that while the Van der Waals state equation (5) is valid, equation (10) is true also.

This completes the formal proof of increasing of the entropy of the Van der Waals gas in the infinitesimal process of the free adiabatic expansion in vacuum. The process with finite change of the volume can be regarded as an integral sum of corresponding infinitesimal processes. Hence, any process of free adiabatic expansion in vacuum for the Van der Waals gas is characterized by the entropy increase. Presupposed violation of thermodynamics second law is apparent since we can not vary parameters in equation (9) independently of one another.

References

- [1] Bazarov I P 1993 *Mistakes and errors in thermodynamics* (Moscow: Moscow University Press)
- [2] Bazarov I P 1964 *Thermodynamics* (Oxford: Pergamon Press)
- [3] Kvasnikov I A 2002 *Thermodynamics and statistical physics* (Moscow: Editorial URSS) vol 1
- [4] Lieb E and Yngvason J 1999 *Physics Report* **310** 1-69
- [5] Kubo R 1968 *Thermodynamics* (Amsterdam: North-Holland Publishing Company) p 170
- [6] Kubo R 1968 *Thermodynamics* (Amsterdam: North-Holland Publishing Company) p 15
- [7] Kubo R 1968 *Thermodynamics* (Amsterdam: North-Holland Publishing Company) p 119